



SKAAPWOLLEK28

CLASS PROTEIN, MINERAL AND TRACE-MINERAL SUPPLEMENT FOR RUMINANTS
REG NR. V19826 ACT 36 OF 1947

BAG CODE ●●●

- A ready-mixed supplement for woolled and mutton sheep and specifically ewes with lambs at foot grazing on dry veld.
- Contains a high energy value.
- Molasses have been added to make this product more palatable and to act as binding agent.
- A specialised lick that stimulates milk production for ewes and promotes high-quality wool growth.
- Contains high levels of natural rumen non-degradable protein as well as sulphur to ensure the amino-acid requirements for improved milk and wool production are met.
- Feed throughout the year, to all sheep classes.
- To stimulate milk production, it should be fed to ewes from four weeks before to at least six weeks after lambing, at between 250g and 500g/animal/day.
- Feed the lick alongside any roughage source.
- An organic barrier system in the lick prevents bloat, especially in maize fields.
- The addition of anion salts to the lick limits the danger of bladder and kidney stones.

COMPOSITION	g/kg
PROTEIN (min)	280
NPN as % of PROTEIN (max)	63%
UREA (max)	50
ENERGY (estimated)	8.5MJ/kg
FIBRE (max)	100
MOISTURE (max)	120
CALCIUM (max)	20
PHOSPHORUS (min)	8
MAGNESIUM (min)	1.64
SULPHUR (min)	7.4
	mg/kg
COPPER (min)	30
MANGANESE (min)	110
ZINC (min)	110
COBALT (min)	0.5
IODINE (min)	1.3
SELENIUM (min)	0.5

RECOMMENDED INTAKES* (g/animal/day)	
Sheep	150 - 450

- WARNING**
- This animal fodder contains an NPN source and should be fed strictly according to instructions. Do not feed together with any other animal fodder containing NPN. Consult a livestock expert.
 - When the transition is made from an NPN-free fodder to a product containing NPN, it is best to feed a 50/50 mixture of the old and the new product for an adjustment period of four to six days.
 - Vinegar works efficiently against NPN poisoning. Mix with equal parts of water, and provide half a bottle per large sheep (1 bottle = 750ml).
 - Keep this fodder out of the rain. Urea is soluble, and animals that drink such a solution can be poisoned.

* Good quality roughage needs to be freely available at all times. The indicated intake only serves as a guideline of the lick amounts that need to be supplied alongside roughage to address nutrient needs. The lick intake can vary depending on roughage quantity and quality.