

**40**

VETMAAKKONSENTRAAT**40**

CLASS PROTEIN, MINERAL AND TRACE-MINERAL SUPPLEMENT FOR RUMINANTS
REG NR. V 21799 ACT 36 OF 1947

BAG CODE ★★

- Ensures optimal intake, digestion and absorption of all necessary nutrients.
- Ensures balanced levels of all nutrients animals need for optimal growth and development.
- Leads to optimal body-mass gain to reach the required marketable target mass within the most economic time.
- Creates a healthy rumen environment that keeps the animals healthy and improves breeding.
- Provide fodder at a ratio of 1% of body mass when animal is placed in the feedlot. Increase every second day with ½% of body mass until maximum fodder intake is obtained.
- Freely provide roughage for the first 10 days in the feedlot. Subsequently withdraw the roughage over a period of time until the animals have sufficiently adapted to their new environment.

COMPOSITION	g/kg
PROTEIN (min)	400
FROM UREA	73.8%
UREA (max)	50
FAT (min)	25
FIBRE (max)	100
MOISTURE (max)	120
CALCIUM (max)	14
PHOSPHORUS (min)	7

WARNING

- This animal fodder contains an NPN source and should be fed strictly according to instructions. Do not feed together with any other animal fodder containing NPN. Consult a livestock expert.
- When the transition is made from an NPN-free fodder to a product containing NPN, it is best to feed a 50/50 mixture of the old and the new product for an adjustment period of four to six days.
- Vinegar works efficiently against NPN poisoning. Mix with equal parts of water, and provide half a bottle per calf or large sheep and two to four bottles per head of cattle (1 bottle = 750ml).
- Keep this product out of the rain. Urea is soluble, and animals that drink such a solution can be poisoned.
- Animals should be given time to adapt. Sufficient pastures and/or roughage should be available at all times.

FEEDING INSTRUCTIONS**

The following are examples of applications of fattening concentrate in ruminant diets.

	Growth feeding		Full rounding off		Full rounding off of sheep		Full rounding off of sheep		Full feed for cattle	
	Bags	kg	Bags	kg	Bags	kg	Bags	kg	Bags	kg
MIXING INSTRUCTIONS										
<i>Vetmaakkonsentraat 40</i>	4	200	3	150	2	100	3	150	2	100
<i>Maize</i>	12	600	12	600	14	700	13	650	12	600
<i>Molasses meal</i>	2	80	3	120	2	80	3	120	–	–
<i>Lucerne meal</i>	–	120	–	–	–	120	–	–	–	–
<i>Roughage</i>	–	–	–	130	–	–	–	80	–	300
COMPOSITION OF MIX	1000		1000		1000		1000		1000	
<i>Protein* (min) g/kg</i>	148		133		140		145		116	
<i>*from urea</i>	*19.39%		*16.18%		*10.18%		*14.18%		*29.69%	
<i>Fat (min)</i>	25		25		25		25		25	
<i>Fibre (max)</i>	100		120		120		120		150	
<i>Moisture (max)</i>	120		120		120		120		120	
<i>Calcium (max)</i>	8.5		10		9		10		6	
<i>Phosphate (min)</i>	4.4		4		3.5		4		3	

** Good quality roughage needs to be freely available at all times. The indicated intake only serves as a guideline of the lick amounts that need to be supplied alongside roughage to address nutrient needs. The lick intake can vary depending on roughage quantity and quality.