



VOLLEDIGESKAAPGROEI

CLASS PROTEIN, MINERAL AND TRACE-MINERAL SUPPLEMENT FOR RUMINANTS
REG NR. V22020 ACT 36 OF 1947

BAG CODE ▲▲▲

- Optimum growth is attained with this feed, but provision needs to be made for sufficient floor space for feed and water, and protection against environmental elements needs to be provided.
- Clean and cool water needs to be provided at a rate of 7ℓ per animal per day.
- This product:
 - contains a safe growth stimulant.
 - has been buffered to protect animals from acidosis.
 - contains anionic salts to limit the occurrence of bladder and kidney stones.

COMPOSITION	g/kg
PROTEIN (min)	140
PROT. EX NPN (max)	26.1%
UREA (maks)	10
FAT (min)	25
FAT (max)	70
FIBRE (min)	110
FIBRE (max)	150
MOISTURE (max)	120
CALCIUM (max)	15
PHOSPHORUS (min)	3

RECOMMENDED INTAKES

This is a complete ration and should be fed ad lib to young animals after an accustoming period of 14 days up until marketing.

Accustoming period: Limit the intake of this finishing feed to approximately 1% of live weight for the first 14 days to prevent feed-related disorders. In addition, provide high-quality roughage ad lib during this period.

WARNING

- This animal fodder contains an NPN source and should be fed strictly according to instructions. Do not feed together with any other animal fodder containing NPN. Consult a livestock expert.
- When an NPN-free product is replaced with a product containing NPN, it is advisable to feed a 50/50 mixture of the old and new product for a transitional period of 4-6 days.
- Vinegar works efficiently against NPN poisoning. Mix with equal parts of water, and provide half a bottle per large sheep (one bottle = 750ml).
- Keep this fodder out of the rain. Urea is soluble, and animals that drink such a solution can be poisoned.