



# WINTERLEK50

CLASS PROTEIN, MINERAL AND TRACE-MINERAL SUPPLEMENT FOR RUMINANTS  
REG NR. VI9830 ACT 36 OF 1947

BAG CODE ■■■

- Cost-effective ready-mixed winter lick.
- Feed to cattle on high-quality winter pastures or maize stubble.
- Contains 50% protein (partially natural protein).
- Feed freely to cattle with access to sufficient dry matter, not where winter pastures are limited or of low quality.
- Prevent lick intakes that are too high.
- Molasses have been included to ensure a dust-free, palatable product.
- The inclusion of a small amount of maize supplies sufficient energy to stimulate the rumen microbes for maximum fibre digestion and NPN utilisation.
- Mix with maize meal to produce production lick.
- Supply supplementation on stubble fields, and aim for the stimulation of intake and the creation of a better environment for the rumen organisms so that effective digestion can take place. Supplementation should not be seen as a substitute for natural pastures.

COMPOSITION	g/kg
PROTEIN (min)	500
NPN as % of PROTEIN (max)	90%
UREA (max)	145
FIBRE (max)	100
MOISTURE (max)	120
CALCIUM (max)	35
PHOSPHORUS (min)	18
MAGNESIUM (min)	1.4
SULPHUR (min)	9
	mg/kg
COPPER (min)	75
MANGANESE (min)	250
ZINC (min)	250
COBALT (min)	0.75
IODINE (min)	3
SELENIUM (min)	1

RECOMMENDED INTAKES* (g/animal/day)	
Cattle	300 - 500
Sheep	Not recommended for sheep

WARNING	
<ul style="list-style-type: none"> <li>• This animal fodder contains an NPN source and should be fed strictly according to instructions. Do not feed together with any other animal fodder containing NPN. Consult a livestock expert.</li> <li>• When the transition is made from an NPN-free fodder to a product containing NPN, it is best to feed a 50/50 mixture of the old and the new product for an adjustment</li> </ul>	<ul style="list-style-type: none"> <li>period of four to six days.</li> <li>• Vinegar works efficiently against NPN poisoning. Mix with equal parts of water, and provide half a bottle per calf and two to four bottles per head of cattle (1 bottle = 750ml).</li> <li>• Keep this fodder out of the rain. Urea is soluble, and animals that drink such a solution can be poisoned.</li> </ul>

\* Good quality roughage needs to be freely available at all times. The indicated intake only serves as a guideline of the lick amounts that need to be supplied alongside roughage to address nutrient needs. The lick intake can vary depending on roughage quantity and quality.