

**P4****BIOFOSP4**

CLASS PROTEIN, MINERAL AND TRACE-MINERAL SUPPLEMENT FOR RUMINANTS
REG NR. V19828 ACT 36 OF 1947

BAG CODE ●●●

- Summer-winter transition lick that contains urea and natural protein.
- The inclusion of a small amount of maize supplies sufficient energy to stimulate the rumen microbes for maximum fibre digestion and NPN utilisation.
- Molasses have been included to make the product dust-free and palatable and to supply energy.
- Ideal product to use as transition lick from summer to winter or as a summer lick on mature summer pastures that have already seeded.

COMPOSITION	g/kg
PROTEIN (min)	200
NPN as % of PROTEIN (max)	79%
UREA (max)	55
FIBRE (max)	100
MOISTURE (max)	120
CALCIUM (max)	80
PHOSPHORUS (min)	40
MAGNESIUM (min)	1.18
SULPHUR (min)	0.7
	mg/kg
COPPER (min)	104
MANGANESE (min)	525
ZINC (min)	525
COBALT (min)	1.6
IODINE (min)	6.25
SELENIUM (min)	1.6
RECOMMENDED INTAKES* (g/animal/day)	
Sheep	35 - 45
Cattle	150 - 250
Lactating cows	200 - 290
WARNING	

- This animal fodder contains an NPN source and should be fed strictly according to instructions. Do not feed together with any other animal fodder containing NPN. Consult a livestock expert.
- When the transition is made from an NPN-free fodder to a product containing NPN, it is best to feed a 50/50 mixture of the old and the new product for an adjustment period of four to six days.
- Vinegar works efficiently against NPN poisoning. Mix with equal parts of water, and provide half a bottle per calf or large sheep and two to four bottles per head of cattle (1 bottle = 750ml).
- Keep this fodder out of the rain. Urea is soluble, and animals that drink such a solution can be poisoned.

* Good quality roughage needs to be freely available at all times. The indicated intake only serves as a guideline of the lick amounts that need to be supplied alongside roughage to address nutrient needs. The lick intake can vary depending on roughage quantity and quality.